

FRANÇOIS SIGRIST

Cyclotomic quadratic forms

Journal de Théorie des Nombres de Bordeaux, tome 12, n° 2 (2000),
p. 519-530

http://www.numdam.org/item?id=JTNB_2000__12_2_519_0

© Université Bordeaux 1, 2000, tous droits réservés.

L'accès aux archives de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Cyclotomic quadratic forms

par FRANÇOIS SIGRIST

RÉSUMÉ. L'algorithme de Voronoï est un procédé permettant d'obtenir la liste complète des formes quadratiques positives parfaites à n variables. Sa généralisation aux G -formes permet de classer les formes G -parfaites, avec l'avantage de se dérouler dans un espace de dimension plus petite (G est un sous-groupe fini de $GL(n, \mathbb{Z})$). On étudie ici la représentation standard du groupe cyclique $G = C_m$ en dimension $\phi(m)$, de polynôme caractéristique $\Phi_m(x)$ (polynôme cyclotomique). Une forme G -invariante est dite forme cyclotomique. Toute les formes G -parfaites sont données pour $\phi(m) < 16$, de même que pour $m = 17$, où la forme cyclotomique la plus dense est entièrement nouvelle. On obtient ainsi une constante d'Hermite cyclotomique, qui s'avère être souvent meilleure que la constante d'Hermite habituelle. C'est le cas pour $m = 5, 7, 11, 13, 16, 17, 36$, et vraisemblablement 32 (les calculs pour $m = 32$ sont en cours, et ont déjà fourni 4600 formes C_{32} -parfaites). Les résultats complets sont disponibles à <http://www.unine.ch/math>.

ABSTRACT. Voronoï's algorithm is a method for obtaining the complete list of perfect n -dimensional quadratic forms. Its generalization to G -forms has the advantage of running in a lower-dimensional space, and furnishes a finite, and complete, classification of G -perfect forms (G is a finite subgroup of $GL(n, \mathbb{Z})$). We study the standard, $\phi(m)$ -dimensional irreducible representation of the cyclic group C_m of order m , and give the, often new, densest G -forms. Perfect cyclotomic forms are completely classified for $\phi(m) < 16$ and for $m = 17$. As a consequence, we obtain precise upper bounds for the Hermite invariant of cyclotomic forms in this range. These bounds are often better than the known or conjectural values of the Hermite constant for the corresponding dimensions; this is indeed the case for $m = 5, 7, 11, 13, 16, 17, 36$. The complete results can be taken from <http://www.unine.ch/math>.

1. Introduction

Denote by $\Phi_m(x) = 1 + c_1x + \dots + c_{n-1}x^{n-1} + x^n$ the m -th cyclotomic polynomial, of degree $n = \phi(m)$. Let $G \in GL(n, \mathbb{Z})$ be the matrix

$$\begin{pmatrix} 0 & 0 & \cdot & \cdot & \cdot & 0 & -1 \\ 1 & 0 & \cdot & \cdot & \cdot & 0 & -c_1 \\ 0 & 1 & \cdot & \cdot & \cdot & 0 & -c_2 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & \cdot & \cdot & \cdot & 1 & -c_{n-1} \end{pmatrix}$$

generating the standard n -dimensional representation of the cyclic group C_m . A *cyclotomic form* is a real quadratic form $q(x) = \sum a_{ij}x_i x_j$ in n variables, invariant under this action of C_m : the symmetric matrix $A = (a_{ij})$ satisfies $G^t A G = A$.

An easy computation shows that the matrix A has then constant diagonals ($a_{i+k,j+k} = a_{i,j}$), and is completely known from the first half of its first row: the space of cyclotomic forms is a linear subspace of $Sym_n(\mathbb{R})$ of dimension $n/2$. Nevertheless, for convenience, we always shall give a cyclotomic matrix by its full first row.

A positive definite real quadratic form has

- a *minimum* $\underline{m} = \min(x^t A x | x \in \mathbb{Z}^n - \{0\})$.
- a set $S = \{\pm v_1, \pm v_2, \dots, \pm v_s\}$ of pairs of *minimal vectors* in \mathbb{Z}^n , satisfying $v_i^t A v_i = \underline{m}$. $2s$ is called the *kissing number* of the form.
- a *Hermite invariant* $\gamma_n(A) = \underline{m} \cdot (\det A)^{-1/n}$.

There is a familiar dictionary linking positive definite forms and Euclidean lattices, via the Gram matrix. A Euclidean lattice has an associated sphere packing, and the packing density δ is related to the Hermite invariant by the relation $\delta = \text{Const}(n) \cdot \gamma_n(A)^{n/2}$: a dense form means a high Hermite invariant.

A quadratic form is *perfect* if it is the unique quadratic form having minimum \underline{m} and set S of minimal vectors. Similarly, a cyclotomic form is said to be C_m -*perfect* if it is the unique *cyclotomic form* with \underline{m} and S as before. A quadratic form is *extreme* if its density is locally maximal in the space of quadratic forms. A cyclotomic form is C_m -*extreme* if its density is locally maximal in the space of *cyclotomic forms*.

An extreme form is always perfect. For the converse, a theorem of Voronoï requires a condition called eutaxy (cf [Mar]). These properties carry over verbatim to the case of C_m -forms [B-M]. A C_m -perfect form is proportional to an integral form, as are the usual perfect forms.

There is a second useful dictionary linking rational cyclotomic forms to the cyclotomic field $\mathbb{Q}(\zeta_m)$. If A is (the matrix of) a cyclotomic form,

there is a unique real element $\alpha \in \mathbb{Q}(\zeta_m)$ such that $x^t Ax = \text{Trace}(\alpha x \bar{x})$. The practical computation runs as follows: write \vec{a} for the first column of the matrix A , and associate to $\alpha = \alpha_0 + \alpha_1 \zeta_m + \dots + \alpha_{n-1} \zeta_m^{n-1}$ the column $\vec{\alpha} = (\alpha_0, \alpha_1, \dots, \alpha_{n-1})^t$. Then $\vec{a} = D \vec{\alpha}$ with matrix $D = (d_{ij}) = \text{Trace}(\zeta_m^{i-j})$. The determinant of D is the discriminant of the field $\mathbb{Q}(\zeta_m)$, and is therefore nonzero: A and α determine each other. If a cyclotomic form is positive definite, the element α is totally positive ($\alpha \gg 0$): all its conjugates are positive. Further, if u is a unit, the cyclotomic form corresponding to $u \cdot \bar{u} \cdot \alpha$ is clearly C_m -equivalent to the original form. This method is extremely useful for finding, or checking, G -equivalences.

Taking determinants in $x^t Ax = \text{Trace}(\alpha x \bar{x})$, one gets (using the fact that α is real):

$$\det(A) = \det(D) \cdot \text{Norm}_{\mathbb{Q}(\zeta_m)/\mathbb{Q}}(\alpha) = \text{Disc}(\mathbb{Q}(\zeta_m)) \cdot (\text{Norm}_{\mathbb{Q}(\zeta_m + \zeta_m^{-1})/\mathbb{Q}}(\alpha))^2.$$

It follows that the determinant of an integral cyclotomic form is equal to $k(m)d^2$, where $k(m)$ is the square-free part of the discriminant of the cyclotomic field $\mathbb{Q}(\zeta_m)$. Further, using the detailed study of decomposition of primes in cyclotomic fields, E. Bayer-Fluckiger and J. Martinet obtained the following useful

Proposition ([Bay-M]). *If q is a prime power dividing d , and prime to m , then $q \equiv \pm 1 \pmod{m}$.*

Up to equivalence, there are only finitely many perfect quadratic forms. Voronoi's algorithm, devised in 1908 [Vor], constitutes a method of explicit complete classification. The densest quadratic form(s) in each dimension can be found with an entirely mechanical computation, the only problem being the gigantic computational complexity.

For cyclotomic forms, it was shown in [B-M-S] that the orthogonal projection of Voronoi's algorithm onto the (dual) space of cyclotomic forms provides an exhaustive classification. Later, it was proved [JCb] that the number of C_m -inequivalent C_m -perfect forms is finite. The practical advantage lies in the dimension of the space of forms ($n/2$ instead of $n(n+1)/2$), bringing the computing time to almost nothing for $n \leq 12$, and to "tolerable" values for the groups C_{17} and C_{32} ($n = 16$). There are 1344 C_{17} -perfect forms, and at least 4600 C_{32} -perfect forms. Both densest forms are completely new.

As is customary in algebraic number theory, we take m to be either odd or divisible by 4.

For many values of m , the conjectured (or proved) densest form (D_4 , E_6 , E_8 , K_{12} , Λ_{16}) appears as a C_m -form. This gives new matrix representations of these well-known forms, and is particularly suitable for further computation of their algebraic invariants.

Even with a long reference list, I cannot do justice to those who helped me during the preparation of this article. However, I wish to thank my

Bordeaux colleagues Anne-Marie Bergé and Jacques Martinet, for their help with many crucial points.

For general results on quadratic forms and lattices, we refer to the books by Conway-Sloane [C-S], and Martinet [Mar].

2. The Voronoï algorithm

Start with a quadratic form $q(x) = x^t A x$, with minimum \underline{m} , and minimal vectors $\{\pm v_i\}$, ($1 \leq i \leq s$). Endow the space $Sym_n(\mathbb{R})$ of symmetric matrices with the Euclidean structure given by $A \cdot B = \text{Trace}(AB)$. Consider the s matrices $V_i = v_i v_i^t$ as elements of the dual space $Sym_n(\mathbb{R})^*$.

The simple observation: $\underline{m} = v_i^t A v_i = \text{Trace}(v_i^t A v_i) = A \cdot V_i$ exhibits the matrix coefficients a_{ij} as a solution of the system of linear equations $A \cdot V_i = \underline{m}$.

The *Voronoï domain* of the matrix A , denoted by $\mathcal{D}(A)$, is the convex hull, in $Sym_n(\mathbb{R})^*$, of the rays λV_i , $\lambda \geq 0$; it is a convex polyhedral cone. Its dimension in $Sym_n(\mathbb{R})^*$ is called the *perfection rank* of A , its codimension the *default of perfection* of A . A perfect form has default zero by definition: the system of equations $X \cdot V_i = \underline{m}$ has $X = A$ as its unique solution. The dual of the Voronoï domain is, by construction, the subset $\{X | X \cdot V_i = v_i^t X v_i \geq 0\}$ of $Sym_n(\mathbb{R})$.

Suppose first that A is not perfect. Then there is a matrix $F \neq 0$ vanishing on all v_i . If $|\theta|$ is small enough, all forms $A(\theta) = A + \theta F$ have same minimum \underline{m} and same set of minimal vectors. There are two values $\rho_1, \rho_2 > 0$ (possibly infinite, but not both) such that $A(-\rho_1)$ and $A(\rho_2)$ still have minimum \underline{m} , but a strictly larger set of minimal vectors. Consequently, after a finite number of steps, one reaches a perfect form.

If A is perfect, the domain $\mathcal{D}(A)$ is of maximal dimension, and is bounded by faces which are hyperplanes in $Sym_n(\mathbb{R})^*$. Given such a face \mathcal{F} , we denote by F the (uniquely defined, up to a positive scalar factor) matrix which is the inner normal to \mathcal{F} .

It is convenient, at this stage, to give the practical method for finding all faces of $\mathcal{D}(A)$. A face matrix F has the following algorithmic description: If $S = \{\pm v_i\}$ is the set of minimal vectors of A , there is a non-trivial partition $S = S_1 \cup S_2$ satisfying:

- $v_i^t F v_i = 0$ for all $v_i \in S_1$ (the V_i lie in the face \mathcal{F}).
- $v_i^t F v_i > 0$ for all $v_i \in S_2$.
- F is unique, up to a positive scalar factor.

The important fact here is that this computation runs entirely in $Sym_n(\mathbb{R})$, though, clearly, the geometric description in $Sym_n(\mathbb{R})^*$ is more convenient for understanding the process.

Take now a perfect form A , F one of its face matrices, and consider the form $A(\theta) = A + \theta F$, with $\theta \geq 0$. If $\theta > 0$ is small enough, $A(\theta)$ still

has minimum \underline{m} , but has default of perfection 1, because, by construction above, its set of minimal vectors is S_1 . Define $\rho = \sup (\theta \mid A(\theta) \text{ has minimum } \underline{m})$. If ρ is infinite, the matrix F is semi-positive definite, and the face \mathcal{F} is said to be a “cul-de-sac”. As was observed in [JCa], and implicitly used in [Vor], this does not happen in the $Sym_n(\mathbb{R})$ - case, but should be kept in mind in generalizing this process. If ρ is finite, $A(\rho)$ is again a perfect form: the *neighbouring form of A along \mathcal{F}* .

The neighbouring relation gives the set of perfect forms the structure of a graph, and Voronoi’s fundamental result [Vor] asserts that this graph is *connected*. Moreover, the $GL(n, \mathbb{Z})$ -equivalence of forms preserves the neighbouring relation, so a quotient graph can be defined. By standard arguments (e.g. Hermite reduction), the quotient graph is *finite*: this is Voronoi’s algorithm.

Besides the finding of all faces of a domain, a time-consuming task, there are three other practical steps needed. We list them for completeness (More on this in [Vor], [JCa], [Mar]):

- Find all minimal vectors.
- Given a face, find the neighbouring form (i.e. find ρ).
- Decide $GL(n, \mathbb{Z})$ -equivalence between two perfect forms.

If G is a finite subgroup of $GL(n, \mathbb{Z})$, it is possible to run Voronoi’s algorithm in the linear subspace \mathcal{T} of G -invariant forms in $Sym_n(\mathbb{R})$. If A is a G -form, its relative Voronoi domain $\mathcal{D}_G(A)$ is then the *orthogonal projection* of $\mathcal{D}(A)$ onto the dual subspace \mathcal{T}^* of $Sym_n(\mathbb{R})^*$. It is proved in [B-M-S] that the resulting neighbourhood graph is *connected*. The finiteness of the quotient graph is more delicate to obtain (G -equivalences have to be taken in the subgroup of $GL(n, \mathbb{Z})$ preserving \mathcal{T}), and has been proved for G -forms in [JCb]. “Culs-de-sac” can appear (see [B-M-S]), and it also happens that the complete graph is already finite. However, for cyclotomic forms, one has the

Proposition. *If G is a \mathbb{Q} -irreducible subgroup of $GL(n, \mathbb{Z})$, there are no “culs-de-sac” in running G -Voronoi’s algorithm.*

Proof. Irreducibility implies that the orbit of any minimal vector spans the whole space. A positive semi-definite form vanishing on a minimal vector must consequently be the zero form. \square

3. Raw data

We shall separate the lists of cyclotomic forms into three families. In family A, the critical form $(D_4, E_6, E_8, K_{12}, \Lambda_{16})$ is cyclotomic. Family B describes C_p -perfect forms, with p a prime. The remaining cases are listed in family C.

For each family, the first table gives successively m , $\phi(m)$, GP = number of C_m -perfect forms, P = number of C_m -perfect forms which are also perfect in the ordinary sense, H=Hermite invariant of the densest form, with its Name, if known. The second table gives first m , $\phi(m)$, First row, then min, det = minimum, resp. determinant of the densest form, in its smallest integral representative, E, F = number of edges, resp. faces, of its Voronoï domain.

Theorem A. *The data for family A are*

m	$\phi(m)$	GP	P	H	Name
8	4	1	1	1.414	D_4
12	4	1	1	1.414	D_4
9	6	2	2	1.665	E_6
15	8	1	1	2.000	E_8
20	8	1	1	2.000	E_8
24	8	1	1	2.000	E_8
21	12	8	8	2.309	K_{12}
28	12	13	7	2.309	K_{12}
40	16	?	?	2.828	Λ_{16}
60	16	?	?	2.828	Λ_{16}

m	$\phi(m)$	First row	min	det	E	F	Name
8	4	(2, -1, 0, 1)	2	4	3	2	D_4
12	4	(2, -1, 1, 0)	2	4	2	2	D_4
9	6	(2, 1, 0, -1, -1, -1)	2	3	4	3	E_6
15	8	(2, 0, 0, 0, 1, -1, -1, 0)	2	1	8	10	E_8
20	8	(2, 0, 0, 1, -1, 0, 1, -1)	2	1	10	10	E_8
24	8	(2, 1, 0, 0, 1, 1, 0, -1)	2	1	9	8	E_8
21	12	(4, 1, -1, -2, 1, 2, 1, -2, -2, -1, 1, 1)	4	729	15	36	K_{12}
28	12	(4, 2, 1, 0, 0, 0, 1, 0, -1, 0, 0, 0)	4	729	21	76	K_{12}
40	16	((4, -1, 0, -1, 0, 2, -2, 1, -2, 1, 0, -1, 2, -2, -2)	4	256	74	?	Λ_{16}
60	16	(4, -1, 0, 2, -1, 2, 0, 0, 2, -1, 2, 1, -2, 2, -1, 0)	4	256	52	?	Λ_{16}

Comment. In the cases with Λ_{16} , the number of edges of the domain makes the finding of the faces practically impossible.

Before giving the data on family B, let's briefly discuss the Craig lattices $A_n^{(r)}$. These famous lattices are thoroughly investigated in [C-S] and [Mar]. Their direct construction as difference lattices gives them matrices (cf.[C-S] p. 433) which are invariant under the n -dimensional representation of C_{n+1}

having characteristic polynomial $1 + x + \dots + x^{n-1} + x^n$. The diagonals are constant, and the first row is filled from left and from right with binomial coefficients:

$$\binom{2r}{r} \quad -\binom{2r}{r+1} \quad \binom{2r}{r+2} \quad \dots \quad \binom{2r}{r+4} \quad -\binom{2r}{r+3} \quad \binom{2r}{r+2}$$

This description is very convenient, but does not show the value of the minimum, whose determination is a difficult combinatorial problem (cf. [Mar] p. 139).

When $n + 1$ is a prime p , the Craig lattices $A_{p-1}^{(r)}$ are *cyclotomic*. In our range of investigation, their minimum is $2r$, by [Ba-Ba], and their density is pretty high. We shall evidently meet these forms below. In high dimensions, Craig lattices are sometimes the densest known (e.g. $p=151$, cf [C-S] p. 43), although it is generally accepted that the C_p -Hermite constant is lower than the absolute constant. Theorem B below shows that, up to a single, and surprising exception, the densest cyclotomic form is a Craig form.

Theorem B. *The data for family B are*

m	$\phi(m)$	GP	P	H	γ_n	Name
5	4	1	1	1.337	1.414	$A_4^{(1)}$
7	6	2	2	1.511	1.665	$A_6^{(2)}$
11	10	5	5	1.948	2.058	$A_{10}^{(2)}$
13	12	25	23	2.106	2.309	$A_{12}^{(2)}$
17	16	1344	1333	2.513	2.828	New

m	$\phi(m)$	First row	min	det	E	F	Name
5	4	(2, -1, 0, 0)	2	5	2	2	$A_4^{(1)}$
7	6	(4, -2, 1, -1, -1, 1)	4	7^3	3	3	$A_6^{(2)}$
11	10	(4, -2, 0, 0, 1, -1, -1, 1, 0, 0)	4	11^3	10	20	$A_{10}^{(2)}$
13	12	(4, -2, 0, 0, 1, -1, -1, 1, 0, 0, 0)	4	13^3	15	84	$A_{12}^{(2)}$
17	16	(6, -3, 2, -1, -2, 2, -3, 2, 0, 0, 2, -3, 2, -2, -1, 2)	6	$17 \cdot 2^{16}$	28	402	New

Comment. The Craig form $A_{16}^{(3)}$ is the densest Craig form, with minimum 6 and determinant 17^5 . Our new form is denser, because $2^4 < 17$. The proportion of perfect forms is high, and will be explained below.

For the last family of forms, with the exceptions of D_8 (for C_{16}) and D_{16} (for C_{32}), there are no otherwise known C_m -perfect forms. The computation for C_{32} is still under way, but the available data suggest that the

densest form has already been found (dense forms tend to have high kissing numbers, and consequently to appear early in the search).

Theorem C. *The data for family C are*

m	$\phi(m)$	GP	P	H	γ_n
16	8	3	1	1.707	2.000
36	12	23	8	2.106	2.309
32	16	> 4500	?	2.446	2.828

m	$\phi(m)$	First row	min	det	E	F
16	8	(8, 2, -4, -3, 0, 3, 4, -2)	8	$2^2 \cdot 241^2$	5	4
36	12	(36, 18, 8, -1, -6, 0, 18, 18, 14, 0, -14, -18)	36	$4463^2 \cdot 5581^2$	6	6
32	16	(12, 5, 0, 3, 4, 3, 0, -4, 0, 4, 0, -3, -4, -3, 0, -5)	12	$2^2 \cdot 31^2 \cdot 5407^2$	21	190

We close this chapter with a few general facts, gathered from our present sample of ca. 6000 C_m -perfect cyclotomic forms. First, in all cases, there is a spanning orbit of minimal vectors, bringing the minimum of the form on the diagonal of the matrix. This seems to be a general fact, but the low dimension, and the triviality of the ideal class group, could contribute to this. Nevertheless, let's formulate this for perfect cyclotomic lattices:

Conjecture 1. *Any perfect cyclotomic lattice has a spanning orbit of minimal vectors.*

Being more pessimistic leads to:

Conjecture 2. *Any perfect cyclotomic lattice is generated by its minimal vectors.*

Another intriguing fact in our sample is

Observation 1. *All perfect cyclotomic forms are even.*

Comment. In the case the prime 2 is unramified in the cyclotomic field, then *any* cyclotomic form, perfect or not, is even¹. In the ramified case, one has to assume C_m -perfection, but the fact is unexplained.

¹This result is due to J. Martinet. The representative element α , being real, has an even trace. When 2 is unramified, this property implies that the corresponding lattice is even.

4. Two detailed samples

We begin with C_{28} , a case appearing in Theorem A. Notations “H”, “E”, “F”, “min”, “det” are as above. “No” is the numbering used by the Voronoï algorithm, and is simply used here to locate neighbours. “s” is the half kissing number, and “i” the default of perfection. Nbr gives the list of neighbours. These 13 forms are listed by decreasing H, beginning with K_{12} .

Theorem A28. *The C_{28} -perfect forms are*

No	H	s	i	E	F	Nbr	min	det
5	2.309	378	0	21	76	$1^6, 2^3, 4^3, 5^{13}, 6^6, 7^9, 8^6, 9^6, 10^6, 11^6, 12^9, 13^3$	4	3^6
7	2.091	126	1	8	9	$1^2, 5^3, 8, 9, 12^2$	12	$29^2 \cdot 1231^2$
1	2.073	154	0	9	10	$2, 3^2, 4, 5^2, 6, 7^2, 8$	6	587^2
12	2.054	112	1	7	8	$5^3, 6, 7^2, 9, 10$	12	39703^2
4	2.045	168	0	9	12	$1^6, 5^6$	4	$2^{6 \cdot 2}$
10	2.007	112	1	7	8	$2, 3, 5^2, 9^2, 11, 12$	8	$2^{12} \cdot 4003^2$
9	2.005	112	1	7	8	$2, 5^2, 7, 10^2, 11, 12$	8	$29^2 \cdot 139^2$
8	1.978	84	0	6	6	$1, 5^2, 7, 11, 13$	10	$3^6 \cdot 617^2$
3	1.978	98	0	6	6	$1^2, 2, 6, 10, 11$	10	$7^2 \cdot 2381^2$
11	1.974	84	1	6	6	$3, 5^2, 8, 9, 10$	16	$3^6 \cdot 10501^2$
6	1.973	98	0	6	6	$1, 2, 3, 5^2, 12$	6	$7^2 \cdot 113^2$
13	1.928	84	7	6	6	$5^3, 8^3$	8	$3^{12} \cdot 7^2$
2	1.915	98	0	6	6	$1, 3, 5, 6, 9, 10$	4	83^2

The next example is C_{13} , contained in Theorem B. The Craig forms appear, and one can observe, contrary to the preceding case, the high proportion of perfect forms.

Theorem B13. *The C_{13} -perfect forms are*

No	H	s	i	E	F	Nbr	min	det	Name
2	2.106	195	0	15	84	$1^6, 3^6, 4^6, 5^6, 6^6, 7^6, 8^6, 9^6, 10^{12}, 11^6, 12^{12}, 13^6$	4	13^3	$A_{12}^{(2)}$
16	2.060	156	0	10	18	$5^6, 9^6, 14^6$	6	13^5	$A_{12}^{(3)}$
14	2.010	78	0	6	6	$3, 4, 10, 11, 15, 16$	28	$13 \cdot 5^4 \cdot 81043^2$	
4	2.005	78	0	6	6	$2, 5, 6, 12^2, 14$	16	$13 \cdot 71527^2$	
11	1.997	78	0	6	6	$2, 3, 5, 12, 14, 15$	24	$13 \cdot 103^2 \cdot 8111^2$	
5	1.996	78	0	6	6	$2, 4, 9, 10, 11, 16$	32	$13 \cdot 4698461^2$	
20	1.994	78	12	6	6	$7, 9, 13, 18, 19, 23$	78	$13 \cdot 53^2 \cdot 79^2 \cdot 236783^2$	
12	1.992	91	0	7	10	$2^2, 4^2, 6, 10, 11, 17, 21, 25$	12	$13 \cdot 13259^2$	

No	H	s	i	E	F	Nbr	min	det	Name
15	1.985	78	0	6	6	3, 6, 7, 11, 14, 18	18	$13 \cdot 154127^2$	
3	1.983	78	0	6	6	7, 9, 11, 13, 14, 15	20	$13 \cdot 103^2 \cdot 2833^2$	
19	1.981	78	0	6	6	7, 13, 17, 18^2 , 20	24	$13 \cdot 877109^2$	
10	1.980	78	0	6	6	2^2 , 5, 12, 13, 14	24	$13 \cdot 879113^2$	
6	1.975	78	0	6	6	2, 4, 7, 12, 15, 17	24	$13 \cdot 891983^2$	
18	1.972	78	0	6	6	7, 15, 17, 19^2 , 20	16	$13^5 \cdot 467^2$	
13	1.964	78	0	6	6	2, 3, 10, 19, 20, 22	42	$13 \cdot 467^2 \cdot 56731^2$	
7	1.963	78	0	6	6	2, 6, 15, 18, 19, 20	14	$13 \cdot 36451^2$	
23	1.949	78	12	6	6	8, 20, 21, 22^2 , 24	26	$13 \cdot 1093^2 \cdot 1429^2$	
22	1.932	78	0	6	6	8, 13, 21, 23^2 , 24	18	$13 \cdot 181193^2$	
17	1.929	78	0	6	6	6, 12, 18, 19, 24, 25	12	$13 \cdot 16067^2$	
21	1.926	78	0	6	6	8, 12, 22, 23, 24^2	24	$13 \cdot 443^2 \cdot 2341^2$	
24	1.926	78	0	6	6	8, 17, 21^2 , 22, 23	16	$13 \cdot 91079^2$	
9	1.901	78	0	6	6	2, 3, 5, 8, 16, 20	12	$13 \cdot 17551^2$	
8	1.889	91	0	7	12	$2^2, 9^2, 21^2, 22^2,$ $23^2, 24^2$	4	$13 \cdot 5^4$	
25	1.865	78	0	6	6	$12^3, 17^3$	4	$13 \cdot 3^6$	
1	1.615	78	0	6	6	2^6	2	13	$A_{12}^{(1)}$

Comment. There are two non-perfect forms in this list, both with a default of perfection of 12, and both with a minimum divisible by 13. Similarly, in the list of 1344 C_{17} -perfect forms, one finds 11 non-perfect forms, all with a default of perfection of 16, and a minimum divisible by 17. This suggests

Observation 2. *The default of perfection of a C_p -perfect form is divisible by $(p - 1)$.*

Observation 3. *The minimum of an integral C_p -perfect, non-perfect C_p -form is divisible by p .*

Observation 2 was first proved by J. Martinet (unpublished). Here is a more general result (proof of my own) settling the claim:

Lemma. *Let A be the matrix of a positive cyclotomic form. Call $\delta(A)$, resp. $\delta_p(A)$ its default of perfection, resp. default of C_p -perfection. Then $\delta(A) - \delta_p(A)$ is divisible by $(p - 1)$.*

Proof. The group C_p acts on $\mathcal{D}(A)$, fixing $\mathcal{D}_p(A)$. As any fixpoint-free rational representation of C_p has dimension divisible by $(p - 1)$, the codimension of $\mathcal{D}_p(A)$ in $\mathcal{D}(A)$ is divisible by $(p - 1)$. Hence $(p(p - 1)/2 - \delta(A)) - ((p - 1)/2 - \delta_p(A))$ is divisible by $(p - 1)$. The lemma follows. □

Observation 3 remains, at present, at the botanical stage. An argument in favour is the fact that the orthogonal projection of $\mathcal{D}(A)$ on $\mathcal{D}_p(A)$ is the average on orbits, i.e. division by p .

5. Final remarks

Our definition of cyclotomic forms is not as general as is used in algebraic number theory, where it means $\text{Trace}(\alpha x \bar{x})$ on an ideal \mathcal{I} of rank one in $\mathbb{Q}(\zeta_m)$. This general description is very convenient, as is illustrated with the Craig forms $A_{p-1}^{(r)}$, which are obtained with $\alpha = \frac{1}{p}$ and $\mathcal{I} = \mathcal{P}^r$ ($\mathcal{P} = (1 - \zeta_p)$). Further, our densest C_{17} -form was identified by J. Martinet: $\alpha = \frac{1}{p}$ and $\mathcal{I} = \mathcal{P}^{-7}\mathcal{Q}$, where $\mathcal{Q}\bar{\mathcal{Q}} = (2)$. It is denser than $A_{16}^{(3)}$ because $2^8 < 17^2$: this ultimately boils down to the fact that 17 is a Fermat prime! Voronoi's algorithm does not fit well in this context, because it runs under a *fixed* representation of the group C_m . In presence of exotic cyclotomic representations, one has to rerun the algorithm entirely, only changing the matrix G generating the representation at the beginning of the program. There is therefore a cyclotomic Hermite constant for each pair of conjugate ideals in the ideal class group of $\mathbb{Q}(\zeta_m)$. Twisting forms with an ideal can only increase the Hermite constant, and looking at the results of [Ba-Ba] on C_{23} , one deduces for example: for the standard cyclotomic representation, the presumably densest form is $A_{22}^{(4)}$ with a Hermite invariant of 2.94...; twisting with one of the exotic representations has the effect of doubling the minimum 2 of $A_{22}^{(1)}$ to 4, giving it an invariant of 3.46... As the presumed γ_{22} is 3.57..., this shows that C_p -cyclotomic forms can become very dense. However, all cases treated in this paper have a trivial ideal class group, and the Hermite invariant of the densest C_m -form can safely be called the *cyclotomic Hermite constant*.

We did not include results on eutaxy in this paper. One reason is that eutaxy is an absolute notion (a cyclotomic form is C_m -eutactic iff it is eutactic). But in view of the very interesting recent results on the classification of eutactic forms (cf [Mar], chap. IX,4), there is no doubt that the study of cyclotomic eutactic forms will prove fruitful.

References

- [Ba-Ba] C. BACHOC, C. BATUT, *Etude algorithmique de réseaux construits avec la forme trace*. J. Exp. Math. **1** (1992), 183–190.
- [Bay-M] E. BAYER-FLUCKIGER, J. MARTINET, *Formes quadratiques liées aux algèbres semi-simples*. J. reine angew. Math. **451** (1994), 51–69.
- [B-M] A.-M. BERGÉ, J. MARTINET, *Réseaux extrêmes pour un groupe d'automorphismes*. Astérisque **198-200** (1992), 41–66.
- [B-M-S] A.-M. BERGÉ, J. MARTINET, F. SIGRIST, *Une généralisation de l'algorithme de Voronoi pour les formes quadratiques*. Astérisque **209** (1992), 137–158.
- [C-S] J.H. CONWAY, N.J.A. SLOANE, *Sphere Packings, Lattices, and Groups*. Springer-Verlag (1992).
- [JCa] D.-O. JAQUET-CHIFFELLE, *Enumération complète des classes de formes parfaites en dimension 7*. Ann. Inst. Fourier **43** (1993), 21–55.
- [JCb] D.-O. JAQUET-CHIFFELLE, *Trois théorèmes de finitude pour les G-formes*. J. Théor. Nombres Bordeaux **7** (1995), 165–176.

- [Mar] J. MARTINET, *Les réseaux parfaits des espaces euclidiens*. Masson (1996).
[Vor] G. VORONOÏ, *Sur quelques propriétés des formes quadratiques positives parfaites*. J. reine angew. Math. **133** (1908), 97–178.

François SIGRIST
Institut de Mathématiques
Université de Neuchâtel
Rue Emile Argand 11
CH-2007 Neuchâtel
Switzerland
E-mail : francois.sigrist@maths.unine.ch